



ELSEVIER

Trends in the land and ocean carbon uptake

Corinne Le Quéré^{1,2}

Only about 45% of the total CO₂ emitted from fossil fuel burning and land use change stayed in the atmosphere on average during the past few decades. The remaining CO₂ was taken up by the carbon reservoirs (the 'CO₂ sinks') in the ocean and on land. The sinks are sensitive to climate and elevated CO₂ levels. Their efficiency in removing CO₂ emissions from the atmosphere is expected to decrease in the future under increasing atmospheric CO₂ because of their response to elevated CO₂ levels, warming and other climate changes. Recent evidence from observations and models suggests that the efficiency of the sinks could have already decreased in the past few decades, but the uncertainties are very large. There is an urgent need for reducing these uncertainties by better monitoring the CO₂ emissions and sinks, and by improving our understanding of the sinks dynamics.

Addresses

¹ School of Environmental Sciences, University of East Anglia, Norwich, NR4 7TJ, UK

² The British Antarctic Survey, High Cross, Madingley Road, Cambridge, CB3 0ET, UK

Corresponding author: Le Quéré, Corinne (C.Lequere@uea.ac.uk)

Current Opinion in Environmental Sustainability 2010, 2:219–224

This review comes from a themed issue on Carbon and nitrogen cycles
Edited by Josep G Canadell

Received 10 February 2010; Accepted 1 June 2010
Available online 8th July 2010

1877-3435/\$ – see front matter
© 2010 Elsevier B.V. All rights reserved.

DOI 10.1016/j.cosust.2010.06.003

Introduction

Despite important advances in the quantification of CO₂ sinks during the past two decades, including their location [1[•]], magnitude [2], driving processes and underlying time-scales [3[•]], important components in the global CO₂ budget remain poorly constrained. CO₂ emissions from fossil fuel combustion are known to an uncertainty of $\sim\pm 6\%$ [4], CO₂ emissions from deforestation and other Land Use Change (LUC) to an uncertainty of $\sim\pm 50\%$ [5^{••}], atmospheric CO₂ growth rate to an uncertainty of $\sim\pm 4\%$ [6], the ocean CO₂ sink to an uncertainty of $\sim\pm 20\%$ [2,3[•],7] and the land CO₂ sink to an uncertainty of $\sim\pm 35\%$ [5^{••}] (Figure 1). The uncertainties reported here are estimates of $\pm 1\sigma$ assessed directly from observations for atmospheric CO₂ growth rate, and indirectly by expert assessment for

the other components based on the spatial coverage in the data and errors in methods, as in [3[•]].


Relatively little attention had been given to the quantification of the trends in CO₂ emissions and sinks until after the publication of the fourth assessment of the Intergovernmental Panel on Climate Change (IPCC-2007). At the close of IPCC-2007, many questions remained open, particularly regarding the response of the CO₂ sinks to elevated CO₂ and climate change [8]. This paper reviews the recent publications that analyse evidence of recent trends in the CO₂ sinks published since IPCC-2007. It discusses the available evidence and the various interpretations of the underlying drivers, and recommends ways to improve our knowledge and understanding.

Trends in the airborne fraction

The fraction of total CO₂ emissions that remain in the atmosphere – the 'airborne fraction (AF)' – has implications for radiative forcing and global warming. It is thus an important quantity to understand and monitor. By definition, AF is a measure of the capacity of the CO₂ sinks to keep up with the CO₂ emissions, which I call the 'sink efficiency' in this paper. Although AF is only an indirect measure of the sink efficiency, it is the only available long time series based on observations. There are many factors that influence AF: First the rate of increase in CO₂ emissions plays a predominant role because the sinks are limited by processes that have specific time-scales, such as the growth of vegetation on land and the vertical transport of carbon in the ocean. Second, the level of CO₂ in the atmosphere influences the uptake by the sinks because CO₂ fertilisation on land (a main cause of the land sink) should saturate at high CO₂ and the dissolution of CO₂ in the ocean reduces the chemical capacity of the ocean to take up additional CO₂. Finally, the land and/or ocean CO₂ sinks or carbon reservoirs could respond to changes in climate [9,10[•]] and other environmental changes such as ocean acidification.

AF is highly variable because the growth of CO₂ in the atmosphere is strongly influenced by the impact of climate variability on the CO₂ sinks (Figure 2). By contrast, the emissions of CO₂ have generally low year-to-year variability. To detect trends in AF, the known interannual variability induced by El Niño events and volcanic eruptions needs to be removed from the atmospheric CO₂ growth rate. This was done by Canadell *et al.* and Raupach *et al.* [11[•],12] who detected a positive trend in airborne

Figure 1



The global budget of CO₂ for the 1990–2000 (blue) and 2000–2008 (red) based on data in [5**]. Emissions from fossil fuel and land use change are based on economic and deforestation statistics [4,49]. Atmospheric CO₂ growth is measured directly [6]. The land and ocean CO₂ sinks are estimated using observations for the 1990–2000 [3*]. For 2000–2008, the ocean CO₂ sink is estimated using an average of several models, while the land CO₂ sink is estimated from the balance of the other terms. Figure reproduced from [50].


fraction of $0.25 \pm 0.21\%$ per year since 1959, with 90% probability ($p = 0.9$) that the signal exceeds the noise in the data. These studies suggested that while about 40% of the emissions of CO₂ remained in the atmosphere in the 1960s, this fraction increased to about 45% in the past decade (Figure 2).

However, the significance of a trend in AF is also influenced by the uncertainty in CO₂ emissions, particularly the large LUC uncertainty [13–15], which was not directly considered in [11*,12]. In a follow up study, Le Quéré *et al.* [5**] used a set of nine published LUC estimates to quantify the impact of LUC uncertainty on

the trend in AF. The positive trend in AF was found to be robust to the LUC estimates considered. Using a wider range of LUC uncertainty with no prior constraints on LUC, Knorr [16] found that the trend in airborne fraction was positive during 1961–2007 but not significant. Further differences in methodology (e.g. the use of monthly data in [16] versus annual data in [5**]) can explain some of the difference in significance between the two studies, but it is nevertheless clear that the trend in AF is at the detection limit and thus that choices of methodology and prior uncertainties have an important impact on the level of significance of the trend detected.

Independent insight on the trend in AF can be obtained from an examination of its potential origin, by looking at trends in the oceanic and land CO₂ sinks separately.

Figure 2



Fraction of the total emissions of CO₂ that remains in the atmosphere (the airborne fraction – top), fraction taken up by the land (middle) and by the ocean (bottom) Units of GtC/y (billion tons of C per year). The total emissions include fossil fuel burning, cement production (about 3% of emissions), deforestation and other land use change [4,49]. The airborne fraction is calculated from the measured growth in atmospheric CO₂ [6]. The land and ocean fraction are an average (full line) of five and four models, respectively (as in [5**]). The grey shading encompasses the range of model estimates and the uncertainty in emissions. The green line in the middle panel is the fraction of the emissions taken up by the land when the land sink is calculated from the difference of the other terms (land CO₂ sink = total CO₂ emissions – atmospheric growth – ocean sink). The difference between the black and green lines on the middle panel provide a measure of the remaining uncertainty in the global CO₂ budget, which can be primarily attributed to uncertainties in the estimates of land CO₂ fluxes [5**]. Figure updated from [5**].

Trends in the ocean carbon uptake

The oceanic sink of CO₂ is mainly a physico-chemical response of the ocean to increasing atmospheric CO₂. CO₂ dissolves in surface waters and is transported from the surface to the deep ocean by ocean currents. The rate at which carbon is transported to the deep ocean is the main factor limiting the uptake of CO₂ by the ocean. The chemical capacity of the ocean and the response of the ocean to surface warming and other climate changes can further alter the rate of uptake of CO₂ by the oceans.

Repeated surveys of the surface ocean CO₂ have been done in many regions for decades. The completion (and continuation) of repeated surveys has led to the publication of several key papers identifying recent trends in surface ocean CO₂. In large regions of the North Atlantic, oceanic CO₂ has increased at the same rate or faster than atmospheric CO₂ since about 1990 [17–20], indicating a local weakening of the CO₂ sink. A similar signal was detected in parts of the Southern Ocean [21,22] first identified from the spatio-temporal variability of atmospheric CO₂ around Antarctica and elsewhere [23], and in the western equatorial Pacific ([24] as updated in [5**]). By contrast, oceanic CO₂ increased at a slower rate than atmospheric CO₂ in the North Pacific [25], as is expected from a response to increasing atmospheric CO₂ alone [26].

The observed trends in oceanic CO₂ have been attributed to various processes. In the Atlantic, different interpretations have been proposed, including changes in marine productivity [18], surface warming [19], changes in physical circulation [20] and natural variability [27]. In the Southern Ocean, models suggest that the ocean is responding to an increase in surface winds [23,28–30], which has been attributed to stratospheric ozone depletion [31]. However this process is not fully coherent with observations of physical changes in the ocean [32] and the mechanisms by which the oceans respond to enhanced winds remain an active area of research [33]. Models have been used to quantify the impact of changes in climate on the global ocean CO₂ sink. All models used so far show a weakening of the global CO₂ sink when the observed changes in climate are imposed on the models [26,28,5**] (Figure 2). The response further analysed in one model was mainly caused by wind changes in the tropical and Southern ocean and surface ocean warming, with important non-linear effects [26].

Despite the short observational record, recent data and modelling studies suggest that the ocean is responding to recent climate change and variability. However, future projections will be difficult to make without a better knowledge of the underlying processes and of the projected changes in surface wind and atmospheric conditions [34*,35,36].

Trends in the land carbon uptake

The land sink of CO₂ is mainly caused by the fertilisation effect of atmospheric CO₂ on terrestrial vegetation and by the regrowth of forests [3^{*}]. Changes in the land CO₂ sink at the regional or global scale are difficult to detect directly from observations because vegetation and soil biomass are very heterogeneous. Traditionally, the trend in the global land CO₂ sink was estimated by subtracting all other terms in the global CO₂ budget (e.g. land CO₂ sink = total CO₂ emissions – atmospheric growth – ocean sink). However this equation transfers all uncertainties to the land CO₂ sink, and does not test our understanding of the underlying processes.

Understanding and modelling of CO₂ fluxes from terrestrial vegetation have made important progress in recent years. New field measurements, flux towers and forest inventories now provide a better picture of the carbon balance of terrestrial vegetation and the distribution of carbon among compartments [37]. Satellite products such as the leaf area index and fire counts provide constraints on the extent and timing of relevant processes. As a result, improved knowledge exists, for instance, in quantifying the contributions of fires [38,39^{*}] and nitrogen limitation [40–42] on terrestrial fluxes. Improved information is also beginning to emerge in our understanding of the role of diffuse light following volcanic activities [43] and the effect of surface ozone levels on leaf openings [44].

It remains challenging to scale up the existing knowledge on processes to global land CO₂ sinks estimates. Thus the global net CO₂ fluxes estimated by models are still poorly constrained and there remain large differences between models (e.g. [39^{*},45]). Nevertheless several important points emerge from recent model analysis: models can account for ~30% of the year-to-year variance in atmospheric CO₂ [5^{**}], they simulate a larger sink in the 1990s compared to the 1980s in agreement with the CO₂ budget approach [39^{*}], and they allocate the sink half in the tropics (compensating deforestation) and half in the Northern Hemisphere in agreement with atmospheric inverse studies [1^{*}].

Attribution of recent trends

Ocean and terrestrial carbon cycle measurements have limited temporal and spatial coverage, and cannot be used directly to infer trends in the CO₂ sinks at the global scale. Four terrestrial and three ocean models were used by Le Quére *et al.* [5^{**}] to assess the effect of recent climate change and variability on the trends in CO₂ sinks and the AF since the early 1960s. When the models were forced by increasing atmospheric CO₂ alone (no changes in climate), AF of the combined results decreased with time by 0.8% per year. In these models, the time scale regulating the absorption of CO₂ by the sinks was faster than the growth in CO₂ emissions from fossil fuel burning and land use change. However when the models were forced by

both increasing atmospheric CO₂ and changes in climate, AF increased by 0.1% per year, close to the trend estimated from observations. The land and ocean models both contributed to the reversal of the trend in about equal proportions. Climate variability could play an important role in setting the trends of the past few decades. For instance, the 1970s was a particularly warm and wet decade, which led to an enhanced land CO₂ sink early in the time series [39^{*}].

The attribution of a trend in AF to the response of CO₂ sinks to climate change and variability in these models is critically dependent on the ability of the models to correctly characterise the time response of the sinks to increasing atmospheric CO₂ alone. In a discussion paper, Gloor *et al.* [46] argues that it is not possible to attribute trends in AF to the climatic impact on CO₂ sinks because of the dominance of the growth rates of atmospheric CO₂. An in-depth evaluation of global models in the context of AF trend attribution remains to be done.

Although the detection of the AF trend [16] and its attribution [46] have been individually challenged, I argue here as in [5^{**}] that the combined evidence from global ocean and land models, ocean observations and from the AF trend based on observations is coherent enough to maintain that a positive trend in AF in the past decades was 'likely'. Likely here is used following the IPCC definition that the probability of occurrence of a positive trend is greater than 66%. Global models suggest that the positive trend in AF can be explained by the response of the CO₂ sinks to climate change and variability. I think that the level of confidence ascribed is a fair representation of recent evidence. I suggest below ways to reduce uncertainty and improve confidence.

Conclusion

It is only recently that sufficient observations exist to constrain the trends in various components of the global carbon cycle. The detection of trends is thus necessarily controversial, because we are at the detection limit of the information. Another decade of observations will certainly help clarify the significance and causes of the observed trends, but time can be gained by improving synergies between models and observations. Improved constraints on the CO₂ sinks would put us in a far better position to project the expected changes this century, to assess the potential for large and/or abrupt changes in biogeochemical cycles with consequences for future climate, and possibly to even provide independent information to constrain the reported emissions of CO₂ at the level of large regions [47]. Such information is important to sustain climate policies and monitor their implementation.

To produce information relevant to climate policies however, there is an urgent need to reduce uncertainties in recent trends and to improve confidence in their attribu-

tion and in future projections. Progress can be made immediately on three fronts: First the large interannual variability in AF estimated from observations (Figure 2) could be reduced by developing more sophisticated methods to remove known variability, as was done by Thompson *et al.* [48] to remove noise in global temperature estimates. Second, the uncertainty in LUC could be better constrained by focusing separately on the mean LUC and its evolution in time since the early 1960s. A more extensive comparison of LUC from methods using different input data and assumptions would help to quantify a time-dependent uncertainty in LUC, and provide more relevant prior information than was used by either Le Quéré *et al.* [5**] or Knorr [16]. Finally, the attribution of recent trends in AF to underlying processes could be improved by an extensive validation of the trends in CO₂ sinks in models, particularly at the regional scale where *in situ* observations exist. It is thus paramount that repeated observations that can constrain CO₂ trends in models be maintained and expanded in order to improve the reliability of model projections.

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Stephens BB *et al.*: **Weak northern and strong tropical land carbon uptake from vertical profiles of atmospheric CO₂**. *Science* 2007, **316**:1732-1735.

A model-data intercomparison showing that models which best match atmospheric observations produce a strong sink for CO₂ in the tropics, compensating deforestation.

2. Manning AC, Keeling RF: **Global oceanic and land biotic carbon sinks from the Scripps atmospheric oxygen flask sampling network**. *Tellus* 2006, **58B**:95-116.
3. Denman K *et al.*: **Couplings between changes and biogeochemistry**. In *Climate Change 2007: The Physical Science Basis. Contribution of Working Group Report of the Intergovernmental Panel on Climate Change*. Edited by Solomon S, Qin D, Manning M, Chen Z, Marquis M, Miller HL. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press; 2007.

A comprehensive synthesis of knowledge on processes driving variability and trends in the land and ocean CO₂ sinks.

4. Marland G: **Uncertainties in accounting for CO₂ from fossil fuels**. *J Ind Ecol* 2008, **12**:136-139.
5. Le Quéré C *et al.*: **Trends in the sources and sinks of carbon dioxide**. *Nat Geosci* 2009, **2**:831-836.
6. Conway TJ, Tans PP, Waterman LS, Thoning KW, Kitzis DR, Masarie KA, Zhang N: **Evidence of interannual variability of the carbon cycle from the NOAA/CMDL global air sampling network**. *J Geophys Res* 1994, **99**:22831-22855.

7. Gruber N *et al.*: **Oceanic sources, sinks, and transport of atmospheric CO₂**. *Global Biogeochem Cycles* 2009, **23**:GB1005.

8. Meehl K *et al.*: **Global climate projections**. In *Climate Change 2007: The Physical Science Basis. Contribution of Working Group Report of the Intergovernmental Panel on Climate Change*. Edited by Solomon S, Qin D, Manning M, Chen Z, Marquis M, Miller HL. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press; 2007.

9. Frank DC, Esper J, Raible CC, Büntgen U, Trouet V, Stocker B, Joos F: **Ensemble reconstruction constraints on the global carbon cycle sensitivity to climate**. *Nature* 2010, **463**:527-532.

10. Friedlingstein P *et al.*: **Climate-carbon cycle feedback analysis: results from the C4MIP model intercomparison**. *J Climate* 2006, **19**:3337-3353.

A comparison of 11 coupled carbon-climate models that quantifies feedbacks between climate and the global carbon cycle.

11. Canadell JP, Le Quéré C, Raupach MR, Field CB, Buitenhuis ET, Ciais P, Conway TJ, Gillett NP, Houghton RA, Marland G: **Contributions to accelerating atmospheric CO₂ growth from economic activity, carbon intensity, and efficiency of natural sinks**. *Proc Natl Acad Sci U S A* 2007, **104**:18866-18870.

The first synthesis of the global annual CO₂ budget that identifies recent trends in the various components and discusses their driving forces and uncertainty.

12. Raupach MR, Canadell JG, Le Quéré C: **Anthropogenic and biophysical contributions to increasing atmospheric CO₂ growth rate and airborne fraction**. *Biogeosciences* 2008, **5**:1601-1613.

13. Ramankutty N, Gibbs HK, Achard F, DeFries R, Foley JA, Houghton RA: **Challenges to estimating carbon emissions from tropical deforestation**. *Global Change Biol* 2007, **13**:51-66.

14. Ito A *et al.*: **Can we reconcile differences in estimates of carbon fluxes from land-use change and forestry for the 1990s?** *Atmos Chem Phys* 2008, **8**:3291-3310.

15. Grainger A: **Difficulties in tracking the long-term global trend in tropical forest area**. *Proc Natl Acad Sci U S A* 2008, **105**:818.

16. Knorr K: **Is the airborne fraction of anthropogenic CO₂ increasing?** *Geophys Res Lett* 2009, **36**:L21710.

17. Schuster U, Watson AJA: **Variable and decreasing sink for atmospheric CO₂ in the North Atlantic**. *J Geophys Res* 2007, **112**: doi: 10.1029/2006JC003941.

18. Lefèvre N, Watson AJ, Olsen A, Ríos AF, Pérez FF, Johannessen T: **A decrease in the sink for atmospheric CO₂ in the North Atlantic**. *Geophys Res Lett* 2004, **31**: doi: 10.1029/2003GL018957.

19. Corbière A, Metzl N, Reverdin G, Brunet C, Takahashi T: **Interannual and decadal variability of the oceanic carbon sink in the North Atlantic subpolar gyre**. *Tellus B* 2007, **59**:168-179.

20. Schuster U, Watson AJ, Bates NR, Corbière A, Gonzalez-Davila M, Metzl N, Pierrot D, Santana-Casiano M: **Trends in the North Atlantic sea-surface fCO₂ from 1990 to 2006**. *Deep Sea Res* 2009, **56**:620-629.

21. Metzl N: **Decadal increase of oceanic carbon dioxide in Southern Indian surface ocean waters (1991-2007)**. *Deep Sea Res II* 2009 doi: 10.1016/j.dsr2.2009.01.002.

22. Takahashi T *et al.*: **Climatological mean and decadal changes in surface ocean pCO₂, and net sea-air CO₂ flux over the global oceans**. *Deep Sea Res II* 2009 doi: 10.1016/j.dsr2.2008.12.009.

23. Le Quéré C *et al.*: **Saturation of the Southern Ocean CO₂ sink due to recent climate change**. *Science* 2007, **316**:1735-1738.

24. Feely RA, Takahashi T, Wanninkhof R, McPhaden MJ, Cosca CE, Sutherland SC, Carr M-E: **Decadal variability of the air-sea CO₂ fluxes in the equatorial Pacific Ocean**. *J Geophys Res* 2006, **111**:C07S03.

25. Takahashi T, Sutherland SC, Feely RA, Wanninkhof R: **Decadal change of the surface water pCO₂ in the North Pacific: a synthesis of 35 years of observations**. *J Geophys Res* 2006, **111**: doi: 10.1029/2005JC003074.

26. Le Quéré C, Takahashi T, Buitenhuis ET, Rödenbeck C, Sutherland SC: **Impact of climate change on the global oceanic sink of CO₂**. *Global Biogeochem Cycles*; In Press.

27. Thomas H, Friederike Prowe AE, Lima ID, Doney SC, Wanninkhof R, Greenbatch RJ, Schuster U, Corbière A: **Changes in the North Atlantic Oscillation influence CO₂ uptake in the North Atlantic over the past 2 decades**. *Global Biogeochem Cycles* 2008, **22**:GB4027.

28. Wetzel P, Winguth A, Maier-Reimer E: **Sea-to-air CO₂ flux from 1948 to 2003: a model study**. *Global Biogeochem Cycles* 2005, **19**:GB2005.
29. Lovenduski N, Gruber N, Doney SC: **Towards a mechanistic understanding of the decadal trends in the Southern Ocean carbon sink**. *Global Biogeochem Cycles* 2008, **22**:BG3016.
30. Lenton A, Matear R: **Role of the Southern Annular Mode (SAM) in Southern Ocean CO₂ uptake**. *Global Biogeochem Cycles* 2007, **21**:GB2016.
31. Thompson DWJ, Solomon S: **Interpretation of recent Southern hemisphere climate change**. *Science* 2002, **296**:895-899.
32. Böning CW, Dispert A, Visbeck M, Rintoul S, Schwarzkopf SF: **The response of the Antarctic Circumpolar Current to recent climate change**. *Nat Geosci* 2008, **1**:864-869.
33. Sallée JB, Speer KG, Rintoul SR: **Zonally asymmetric response of the Southern Ocean mixed-layer depth to the Southern Annular Mode**. *Nat Geosci* 2010, **3**:273-279.
34. Lenton A, Codron F, Bopp L, Metzl N, Cadule P, Tagliabue A, Le Sommer J: **Stratospheric ozone depletion reduces ocean carbon uptake and enhances ocean acidification**. *Geophys Res Lett* 2009, **36**:L12606.
- A model quantification of the possible future evolution of the Southern Ocean CO₂ sink, taking into account the evolution of climate and the recovery of stratospheric ozone depletion.
35. Zickfeld K, Fyfe JC, Eby M, Weaver AJ: **Comment on "Saturation of the Southern ocean CO₂ sink due to recent climate change"**. *Science* 2008, **319**:570b.
36. Le Quéré C *et al.*: **Response to comments on "Saturation of the Southern Ocean CO₂ sink due to recent climate change"**. *Science* 2008, **319**:570c.
37. Friend AD, Arneth A, Kiang NY, Lomas M, Ogee J, Rodenbeckk C, Running SW, Santaren JD, Sitch S, Viovy N *et al.*: **FLUXNET and modelling the global carbon cycle**. *Global Change Biol* 2007, **13**:610-633.
38. van der Werf GR, Randerson JT, Giglio L, Collatz GJ, Kasibhatla PS, Arellano AF Jr: **Interannual variability in global biomass burning emissions from 1997 to 2004**. *Atmos Chem Phys* 2006, **6**:3423.
39. Sitch S *et al.*: **Evaluation of the terrestrial carbon cycle, future plant geography and climate-carbon cycle feedbacks using five Dynamic Global Vegetation Models (DGVMs)**. *Global Change Biol* 2008, **14**:2015-2039.
- A comparison of four global vegetation models that identifies and discusses the response of terrestrial vegetation to recent CO₂ increase and climate variability.
40. Jain A, Yang X, Kheshgi H, McGuire AD, Post W, Kicklighter D: **Nitrogen attenuation of terrestrial carbon cycle response to global environmental factors**. *Global Biogeochem Cycles* 2009, **23**:GB4028.
41. Zaehle S, Friend AD, Dentener F, Friedlingstein P, Peylin P, Schulz M: **Carbon and nitrogen cycle dynamics in the O-CN land surface model. 2: The role of the nitrogen cycle in the historical terrestrial carbon balance**. *Global Biogeochem Cycles* 2010, **24**:GB 1006.
42. Thornton PE, Lamarque J-F, Rosenbloom NA, Mahowald NM: **Influence of carbon-nitrogen cycle coupling on land model response to CO₂ fertilization and climate variability**. *Global Biogeochem Cycles* 2007, **21**:GB4018.
43. Mercado LM, Bellouin N, Sitch S, Boucher O, Huntingford C, Wild M, Cox PM: **Impact of changes in diffuse radiation on the global land carbon sink**. *Nature* 2009, **458(7241)**:1014-1017.
44. Sitch S, Cox PM, Collins WJ, Huntingford C: **Indirect radiative forcing of climate change through ozone effects on the land-carbon sink**. *Nature* 2007 doi: [10.1038/nature06059](https://doi.org/10.1038/nature06059).
45. Randerson JT *et al.*: **Systematic assessment of terrestrial biogeochemistry in coupled climate-carbon models**. *Global Change Biol* 2009, **15**:2462-2484.
46. Gloor M, Sarmiento JL, Gruber N: **What can be learned about carbon cycle climate feedbacks from CO₂ airborne fraction?** *Atmos Chem Phys Discuss* 2010, **10**:9045-9075.
47. Durant AJ, Le Quéré C, Hope C, Friend A: **Constraints on annual emissions and sinks of carbon dioxide**. *Proc R Soc Lond* 2010; Submitted for Publication.
48. Thompson DWJ, Kennedy JJ, Wallace JM, Jones PD: **A large discontinuity in the mid-twentieth century in observed global-mean surface temperature**. *Nature* 2008, **453**:646-650.
49. Houghton RA: **Revised estimates of the annual net flux of carbon to the atmosphere from changes in land use and land management 1850-2000**. *Tellus* 2003, **55**:378.
50. Le Quéré C: **Closing the global budget for CO₂**. *Global Change* 2009, **74**:28-31.